



## Dompracid P

### Dompracid P

#### Introduction

This tablet belongs to the class of drugs known as Dopamine antagonists along with proton pump inhibitors (PPI) in combination.

#### Description

Dompracid works by tightening the muscles near the entering area of the abdomen and relaxes the muscles present in the exit area of the stomach. This process increases the speed of digestion of food that we intake leading to a reduction in feeling the sickness and nausea. This drug can also stop or decrease stimulation in the 'vomiting center' of the brain which leads to a reduction in the feeling of vomiting and nausea.

#### Uses and Benefits of Dompracid P

The most common use of this medication is:

- Nausea Or Vomiting
- Acidity
- Heartburn
- Gastric Ulcer

#### Composition and Mechanism Of Dompracid P

Dompracid P Tablet attaches to the dopaminergic receptors without causing any release of the chemical dopamine. This, in turn, facilitates gastric emptying and decreases small bowel transit time. Also, it binds to H<sup>+</sup>/K<sup>+</sup>-exchanging ATPase in gastric parietal cells, resulting in blockage of acid secretion. It inhibits the acid secretion in the stomach by inhibiting H<sup>+</sup>/K<sup>+</sup> ATPase channels.

Domperidone: it increases the movements or contractions of the stomach and bowel. Domperidone is also used to treat nausea and vomiting caused by other drugs

Pantoprazole: It belongs to a class of drugs known as proton pump inhibitors (PPIs). Pantoprazole is used to treat certain stomach and esophagus problems (such as acid reflux). It works by decreasing the amount of acid your stomach makes. It helps heal acid damage to the stomach and esophagus, helps prevent ulcers, and may help prevent cancer of the esophagus.

## How to take Dompracid P

- Dompracid P should be swallowed whole.
- Dompracid P should not be crushed, chewed or broken before consumption.
- The dosage prescribed by the doctor should be strictly followed.

## Dompracid P Interaction

- **Interaction with Disease**  
There are no interactions with any disease
- **Interaction with Alcohol**  
There are no interactions with alcohol
- **Interaction with Food**  
There are no interactions with any food.
- **Interaction with Medicine**  
There are no interactions with any medication

There may be other interactions with Dompracid P than those already mentioned above. The interactions also depend on the medical condition of the individual, his/her medical history and the current parallel running medications.

## Coxinip-TH Contraindications:

- Pediatric use: Coxinip-TH should not be given to children below 16 years of age.
- Infections: A patient suffering from any kind of infection should exercise caution while taking Coxinip-TH as it may hide the respective symptoms. It is advised to monitor the condition and a doctor should be consulted.
- Pregnancy: Coxinip-TH is not recommended for pregnant women unless necessary. All the risks and benefits should be calculated and medicine should be consumed only if the benefits outweigh the risks. Coxinip-TH can also be replaced with a suitable substitute if needed.
- Breast-feeding: Coxinip-TH should not be used by breastfeeding women unless necessary. All the risks and benefits should be calculated, and it should be taken only if benefits outweigh the risk. Also, breastfeeding should be discontinued while consuming Dompracid

## Side Effects of Dompracid P

Heart rhythm disorders, comprise a urinary frequency, severe itching of the skin, pain in a group of muscles, may also cause rash, body temperature, diarrhea, blurred vision, dizziness, and headache.

There may be other side effects to Dompracid P than those already mentioned above. The side effects also depend on the medical condition of the individual, his/her medical history and the current parallel running medications.

## Missed Dose

If you are taking this product on a prescribed schedule and miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

**Storage**

Store at room temperature between 59-86 degrees F (15-30 degrees C) away from light and moisture. Do not store in the bathroom. Keep all medicines away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company for more details about how to safely discard your products\



CONCEPTIVE FARMACIA  
ERA OF BIO-MEDICAL REVOLUTION



CONCEPTIVE FARMACIA  
ERA OF BIO-MEDICAL REVOLUTION